

疾病予防控制局

中国 ARI 定点監視状況 (2026 年第 7 週)

疾病予防控制局 www.nhc.gov.cn 2026-02-19

来源: 疾病予防控制局

2026 年第 7 週 (2026 年 2 月 9 日～2 月 15 日)、全国 (香港、マカオ、台湾を除く) の定点病院から採取された外来および救急のインフルエンザ様疾患患者と重症急性呼吸器感染症による入院患者の呼吸器サンプルに対し、新型コロナウイルス、インフルエンザウイルス、RS ウイルス (RSV)、アデノウイルス、ヒトメタニューモウイルス (hMPV)、パラインフルエンザウイルス (PIV)、通常のコロナウイルス、ボカウイルス (HBoV)、ライノウイルス (HRV)、エンテロウイルス (EV) の 10 種類及びマイコプラズマ肺炎 (MPP) といった呼吸器病原体のウイルス検査を実施した。

1. モニタリング結果

第 6 週、全国定点 (哨点) 病院が報告したインフルエンザ様患者の救急外来受診者数の割合 (インフルエンザ様患者の比率) は 3.3%であった; 哨点病院の救急外来のインフルエンザ様症例の呼吸器検体から陽性病率の上位三位の病原体はインフルエンザウイルス (11.0%: 2%の減少)、RSV (8.0%)、PIV (5.4%) であった; 入院患者における重症急性呼吸器感染症の呼吸器検体から検出された陽性病率の上位三位の病原体は RSV (12.8%)、HRV (5.3%)、PIV (5.1%) であった。その他の病原体のモニタリング結果については表 1 を参照のこと。モニタリングの結果、北部と南部で地域や年齢層の間に差異が見られたが、詳細については表 2、表 3 を参照のこと。

2. 分析と健康のためのヒント

全国の発熱外来の診療状況や定点 (哨点) 医院の救急外来におけるインフルエンザ様患者の診療状況や多くの病原体検査等のモニタリング結果を結合して分析すると、検出された病原体はすべて既知の一般的なものであり、未知の病原体やそれらが引き起こす新たな感染症は発見されていない。最近、インフルエンザウイルスと RSV が主な病原体として検出されている。中国全土におけるインフルエンザの流行は減少が続き、あらゆる省で既に中～低レベルに低下しているが、一部の省では通常の流行期レベルにまで減少している。最近の RSV 検査陽性率は南方の省では増加、北方の省では下降という特徴がみられるが、北方の省はその陽性率が南方の省に比して高く、0～4 歳の症例グループにおける RSV の陽性率は他の年齢層よりも高い。PIV の検査陽性率は上昇の傾向にあり、南方の省が北方より高く、0～4 歳の症例グループが他の年齢層より高くなっている。HRV と通常のコロナウイルスの検査陽性率は一定のレベルで増減している。COVID-19 やその他の気道感染症の病原体陽性率は総じて低いレベルにある。

現在、わが国の呼吸器感染症は依然ピーク期にあるが、総じて下降の趨勢にあり、インフルエンザは既に低レベルの流行状態にある。RSV の活動は南北で開きがあり、HRV、PIV 及び一般的なコロナウイルスには一定レベルの活動が見られる。呼吸器感染症の蔓延を防ぐため、以下の保護対策を講じるよう推奨する:

- 科学的なマスクの着用:** 治療中は常にマスク着用を; 人が密集する場所や飛行機、電車、地下鉄などの公共交通機関を利用する際には、マスク着用を推奨するが、特に特に高齢者および一部の慢性疾患のある患者の感染リスクを軽減すること。
- 良好な衛生習慣の維持:** 咳やくしゃみの際は、ティッシュやタオル或いは肘で口と鼻を覆う; 感染リスク減少のため、手指衛生に注意し汚れた手で目や鼻、口に触れないようにすること。
- 健康的なライフスタイルの推進:** バランスの取れた食事、適度な運動、十分な休息をとり、体の免疫力を高めること。発熱や咳などの呼吸器感染症の症状がある場合、人と接触するときはマスクを着用し、室内の換気を良好に保ち、必要に応じて速やかに医師の診察を受けること。
- 積極的なワクチンの接種:** 免疫力が弱い人 (妊婦、幼児、高齢者、慢性疾患患者など) は、予防接種のガイドラインに従い呼吸器感染症のワクチン接種を受けねばならない。

表 1: 呼吸器検体における病原体の核酸陽性検出率 (%) (第 7 週)

病原体	外来インフルエンザ 様疾患	重症急性呼吸器感染症 入院者
-----	------------------	-------------------

	第7周	先週比*	第7周	先週比*
COVID-19	1.7	-0.5	1.1	-0.8
インフルエンザウイルス	11.0	-1.7	4.9	-0.1
RS ウイルス RSV	8.0	-0.5	12.8	+0.1
アデノウイルス Adenovirus	1.9	-0.5	1.2	-0.6
ヒトメタニューモウイルス hMPV	0.4	-0.2	0.6	+0.2
パラインフルエンザウイルス PIV	5.4	-0.1	5.1	+0.7
通常のコロナウイルス Norm. coronavirus	4.6	-0.2	3.0	-1.0
ボカウイルス HBoV	0.4	-0.2	0.8	-0.3
ライノウイルス HRV	5.1	-0.5	5.3	-0.7
エンテロウイルス EV	1.0	-0.3	0.9	+0.2
マイコプラズマ肺炎 MPP	0.2	-0.1	0.3	-0.3

*「+」は今週の特定病原体検出の陽性率が先週と比較して増加したことを、「-」は先週と比較して減少したことを示す。

表2 呼吸器検体における主要病原体の核酸検査陽性率の地域差（第7週）

省	外来インフルエンザ様疾患			重症急性呼吸器感染症入院者		
	第一位	第二位	第三位	第一位	第二位	第三位
南方各省	Influenza virus	PIV	RSV	RSV	PIV	HRV
北方各省	RSV	Influenza virus	HRV	RSV	Influenza virus	HRV

南方の省には、上海、江蘇省、浙江省、安徽省、福建省、江西省、湖北省、湖南省、広東省、広西省、海南省、重慶、四川省、貴州省、雲南省が含まれ、

北方の省には、北京、天津、河北省、山西省、内モンゴル自治区、遼寧省、吉林省、黒竜江省、山東省、河南省、チベット自治区、陝西省、甘肅省、青海省、寧夏回族自治区、新疆ウイグル自治区、新疆生産建設兵団が含まれる。

表3 呼吸器検体における主要病原体の核酸検出陽性率の年齢層別違い（第7週）

年齢グループ	外来インフルエンザ様疾患	重症急性呼吸器感染症入院者
0～4歳	1. RSV	1. RSV
	2. PIV	2. Influenza virus
	3. Influenza virus	3. HRV
5～14歳	1. Influenza virus	1. HRV/ Influenza virus
	2. HRV	2. RSV
	3. RSV	3. Norm. coronavirus/ PIV
15～59歳	1. Influenza virus	1. Norm coronavirus
	2. HRV	2. HRV
	3. Norm. coronavirus	3. RSV
60歳以上	1. Influenza virus	1. RSV
	2. HRV	2. Influenza virus
	3. Norm. coronavirus	3. HRV




図1: 定点病院の外来および救急外来におけるインフルエンザ様症例の割合の週別推移




图 2 定点医院的外来および救急外来におけるインフルエンザ様症例の呼吸器検体における病原体核酸検査陽性率の週別推移




图 3：定点病院における入院患者の重症急性呼吸器感染症症例の呼吸器検体における病原体核酸検査陽性率の週別推移

https://www.chinacdc.cn/jksj/jksj04_14275/202602/t20260219_315150.html

National sentinel surveillance of ARI (week 7, 2026)

China CDC www.nhc.gov.cn 2026-02-19 Source: China CDC

In the week 7 of 2026 (February 9, – February 15, 2026), respiratory specimens from outpatient and emergency influenza-like cases and inpatient severe acute respiratory infection cases collected in sentinel hospitals across the country (excluding Hong Kong, Macao and Taiwan) were tested for novel coronavirus pneumonia, coronavirus, influenza virus, respiratory syncytial virus, adenovirus, human metapneumovirus, parainfluenza virus, common coronavirus, bocavirus, rhinovirus and enterovirus, as well as a variety of respiratory viruses including Mycoplasma pneumoniae Pathogen detection.

1. Monitoring results

In the week 7, the proportion of influenza-like cases reported by sentinel hospitals nationwide to the total number of outpatient visits (percentage of influenza-like cases) was 3.3%; The top three pathogens with positive rates in respiratory samples of influenza-like cases in outpatient and emergency departments of sentinel hospitals were **Influenza virus (11.0%)**, **RSV (8.0%)**, and **PIV (5.4%)**; the top three pathogens with positive rates in respiratory samples of hospitalized severe acute respiratory infection cases were **RSV (12.8%)**, **HRV (5.3%)**, and **PIV (5.1%)**. The monitoring results of other pathogens are shown in Table 1. The monitoring results showed differences between the north and south regions and between different age groups, as shown in Tables 2 and 3.

2. Analysis and health tips

Analysis of nationwide fever clinic treatment data, sentinel hospital outpatient and emergency department influenza-like illness case reports, and multi-pathogen detection results show that all detected pathogens are known

common pathogens, and no unknown pathogens or emerging infectious diseases they cause were found. Recently, **influenza virus** and **RSV** have been the main pathogens detected. The overall influenza epidemic situation nationwide is showing a continuous downward trend, with all provinces having reached medium or low epidemic levels, and some provinces reaching inter-epidemic levels. Recently, the **RSV**'s positivity rate has shown a trend of increasing in southern provinces and decreasing in northern provinces, with the positivity rate still higher in northern provinces than in southern provinces, and higher in the 0–4-year-old case group than in other age groups. The parainfluenza virus positivity rate is showing an upward trend, higher in southern provinces than in northern provinces, and higher in the 0–4-year-old case group than in other age groups. The positivity rates for **HRV** and common coronaviruses fluctuate at a certain level. The positivity rates for COVID-19 and other monitored respiratory pathogens are generally at a low level.

Currently, respiratory infectious diseases are still in their peak season in China, but the overall trend is declining. The national influenza epidemic has dropped to a low prevalence level. RSV activity shows differences between north and south, while HRV, PIV, and common coronaviruses exhibit some activity. To prevent the spread of respiratory infectious diseases, the public is advised to take the following personal protective measures:

- (1) **Wear masks scientifically:** You must wear a mask throughout the medical treatment process; it is recommended to wear a mask in crowded places or when taking public transportation (such as airplanes, trains, subways, etc.), especially for the elderly and some patients with chronic underlying diseases, to reduce the risk of infection.
- (2) **Maintain good hygiene habits:** Cover your mouth and nose with a tissue, towel, or elbow when coughing or sneezing; pay attention to hand hygiene and avoid touching your eyes, nose, and mouth with unclean hands to reduce the risk of pathogen transmission.
- (3) **Advocate a healthy lifestyle:** Maintain a balanced diet, engage in appropriate exercise and get enough rest to enhance the body's immunity. If you experience fever, cough or other respiratory infection symptoms, wear a mask when in contact with others and keep the room well-ventilated; seek medical attention promptly if necessary.
- (4) **Actively vaccinate:** People with weak immunity (such as pregnant women, young children, the elderly and patients with chronic diseases, etc.) should receive relevant vaccines to prevent respiratory infectious diseases according to vaccination guidelines.

Table 1: Positive rate of nucleic acid detection of pathogens in respiratory samples in **Week 7** (%)

Pathogen	Outpatient influenza-like illness		Inpatient severe ARI cases	
	Week 7	Vs Prev week*	Week 7	Vs Prev week*
COVID-19	1.7	-0.5	1.1	-0.8
Influenza virus	11.0	-1.7	4.9	-0.1
Respiratory syncytial virus (RSV)	8.0	-0.5	12.8	+0.1
Adenovirus	1.9	-0.5	1.2	-0.6
hMPV	0.4	-0.2	0.6	+0.2
Parainfluenza virus (PIV)	5.4	-0.1	5.1	+0.7
Norm. coronavirus	4.6	-0.2	3.0	-1.0
Bocavirus (HBoV)	0.4	-0.2	0.8	-0.3
Rhinovirus (HRV)	5.1	-0.5	5.3	-0.7
EV	1.0	-0.3	0.9	+0.2
MPP	0.2	-0.1	0.3	-0.3

Note: "+" indicates that the positive rate of specific pathogen detection this week has increased compared with last week; "-" indicates that the positive rate of specific pathogen detection this week has decreased compared with last week.

Table 2 Regional differences in the positive rates of nucleic acid tests for major pathogens in respiratory samples in **week 7**

Province	Outpatient influenza-like illness			Hospitalized severe acute respiratory infection cases		
	No.1	No.2	No.3	No.1	No.2	No.3
S. Provinces	Influenza virus	PIV	RSV	RSV	PIV	HRV
N. Provinces	RSV	Influenza virus	HRV	RSV	Influenza virus	HRV

The southern provinces include Shanghai, Jiangsu, Zhejiang, Anhui, Fujian, Jiangxi, Hubei, Hunan, Guangdong, Guangxi, Hainan, Chongqing, Sichuan, Guizhou and Yunnan.

The northern provinces include Beijing, Tianjin, Hebei, Shanxi, Inner Mongolia, Liaoning, Jilin, Heilongjiang, Shandong, Henan, Tibet, Shaanxi, Gansu, Qinghai, Ningxia, Xinjiang and Xinjiang Production and Construction Corps.

Table 3 Differences in the positive rate of nucleic acid detection of main pathogens in respiratory samples in week 7 by age group

Age group	Outpatient influenza-like cases	Hospitalized SAR infection cases
0-4 years old	1. RSV	1. RSV
	2. PIV	2. Influenza virus
	3. Influenza virus	3. HRV
5-14 years old	1. Influenza virus	1. HRV/ Influenza virus
	2. HRV	2. RSV
	3. RSV	3. Norm. coronavirus/ PIV
15-59 years old	1. Influenza virus	1. Norm coronavirus
	2. HRV	2. HRV
	3. Norm. coronavirus	3. RSV
60 years old and above	1. Influenza virus	1. RSV
	2. HRV	2. Influenza virus
	3. Norm. coronavirus	3. HRV




Figure 1 Weekly trend of the proportion of influenza-like cases in outpatient and emergency departments of sentinel hospitals to the total number of outpatient and emergency department visits




Figure 2 Weekly trend of the positive rate of nucleic acid test for pathogens in respiratory samples of influenza-like cases in outpatient and emergency departments of sentinel hospitals




Figure 3 Weekly trend of the positive rate of nucleic acid test for pathogens in respiratory samples of inpatient severe acute respiratory infection cases in sentinel hospitals

..... 以下是中国語原文

全国急性呼吸道传染病哨点监测情况（2026 年第 7 周）

疾病预防控制局 www.nhc.gov.cn 2026-02-19

来源：疾病预防控制局

2026 年第 7 周（2026 年 2 月 9 日-2 月 15 日），对全国哨点医院（不含港澳台）采集的门诊急诊流感样病例和住院严重急性呼吸道感染病例的呼吸道样本，开展新型冠状病毒、流感病毒、呼吸道合胞病毒、腺病毒、人偏肺病毒、副流感病毒、普通冠状病毒、博卡病毒、鼻病毒和肠道病毒等 10 种病毒以及肺炎支原体在内的多种呼吸道病原体检测。

一、分析及健康提示

第 7 周，全国哨点医院报告的流感样病例数占门诊急诊就诊总数的比例（流感样病例百分比）为 **3.3%**；哨点医院门诊急诊流感样病例呼吸道样本检测阳性率前三位病原体为**流感病毒（11.0%：2%减少）**、**呼吸道合胞病毒（8.0%）**、**副流感病毒（5.4%）**；住院严重急性呼吸道感染病例呼吸道样本检测阳性率前三位病原体为**呼吸道合胞病毒（12.8%）**、**鼻病毒（5.3%）**、**副流感病毒（5.1%）**。其他病原体监测结果见表 1。监测结果存在南北方地域差异和不同年龄段人群差异，具体见表 2 和表 3。

二、分析及健康提示

结合全国发热门诊（诊室）诊疗情况、哨点医院门诊急诊流感样病例就诊情况、多病原检测等监测结果分析显示，检出均为已知常见病原体，没有发现未知病原体及其导致的新发传染病。近期流感病毒和呼吸道合胞病毒为主要检出病原体。全国流感疫情总体呈持续下降趋势，所有省份已降至中流行水平或低流行水平，个别省份降至流行间期水平。近期呼吸道合胞病毒检测阳性率呈现南方省份上升、北方省份下降的特征，北方省份检测阳性率仍高于南方省份，0~4 岁病例组高于其他年龄组。副流感病毒检测阳性率呈上升趋势，**南方省份高于北方省份，0~4 岁病例组高于其他年龄组**。鼻病毒和**普通冠状病毒**检测阳性率在一定程度波动。新型冠状病毒和其他监测的呼吸道病原体检测阳性率总体处于低水平。

当前，我国呼吸道传染病仍处于高发季节，总体呈下降趋势，**全国流感疫情已降至低流行水平**，呼吸道合胞病毒活动呈现南北差异，鼻病毒、副流感病毒和普通冠状病毒存在一定程度活动。为防范呼吸道传染病疫情，建议公众采取以下个人防护措施：

- （1）科学佩戴口罩：在就医过程中需全程佩戴口罩；在人群密集场所或乘坐公共交通工具（如飞机、火车、地铁等）时建议佩戴口罩，尤其是老年人和一些有慢性基础性疾病的患者，以减少感染风险。
- （2）保持良好卫生习惯：咳嗽或打喷嚏时，用纸巾、毛巾或手肘遮住口鼻；注意手卫生，避免用不洁净的手触摸眼、鼻、口，以减少病原体传播风险。

- (3) 倡导健康生活方式：要均衡饮食、适量运动和充足休息，增强身体免疫力。出现发热、咳嗽或其他呼吸道感染症状时，与人接触要佩戴口罩，保持室内良好通风；根据需要及时就医。
- (4) 积极主动接种疫苗：免疫力较弱人群（如孕妇、低龄儿童、老人及慢性病患者等）应根据疫苗接种指引接种预防呼吸道传染病的相关疫苗。

表 1 第 7 周呼吸道样本病原体核酸检测阳性率（%）

病原体	门急诊流感样病例		住院严重急性呼吸道感染病例	
	第 7 周	较上周*	第 7 周	较上周*
新型冠状病毒	1.7	-0.5	1.1	-0.8
流感病毒	11.0	-1.7	4.9	-0.1
呼吸道合胞病毒	8.0	-0.5	12.8	+0.1
腺病毒	1.9	-0.5	1.2	-0.6
人偏肺病毒	0.4	-0.2	0.6	+0.2
副流感病毒	5.4	-0.1	5.1	+0.7
普通冠状病毒	4.6	-0.2	3.0	-1.0
博卡病毒	0.4	-0.2	0.8	-0.3
鼻病毒	5.1	-0.5	5.3	-0.7
肠道病毒	1.0	-0.3	0.9	+0.2
肺炎支原体	0.2	-0.1	0.3	-0.3

注：“+”表示本周特定病原体检测阳性率数值较上周增加；
“-”表示本周特定病原体检测阳性率数值较上周下降。

表 2 第 7 周呼吸道样本主要病原体核酸检测阳性率区域差异

省份	门急诊流感样病例			住院严重急性呼吸道感染病例		
	第一位	第二位	第三位	第一位	第二位	第三位
南方省份	流感病毒	副流感病毒	呼吸道合胞病毒	呼吸道合胞病毒	副流感病毒	鼻病毒
北方省份	呼吸道合胞病毒	流感病毒	鼻病毒	呼吸道合胞病毒	流感病毒	鼻病毒

南方省份包括：上海、江苏、浙江、安徽、福建、江西、湖北、湖南、广东、广西、海南、重庆、四川、贵州和云南。
北方省份包括：北京、天津、河北、山西、内蒙古、辽宁、吉林、黑龙江、山东、河南、西藏、陕西、甘肃、青海、宁夏、新疆和新疆生产建设兵团。

年龄组	门急诊流感样病例	住院严重急性呼吸道感染病例
0~4 岁	①呼吸道合胞病毒	①呼吸道合胞病毒
	②副流感病毒	②流感病毒
	③流感病毒	③鼻病毒
5~14 岁	①流感病毒	①鼻病毒·流感病毒
	②鼻病毒	②呼吸道合胞病毒
	③呼吸道合胞病毒	③普通冠状病毒·副流感病毒
15~59 岁	①流感病毒	①普通冠状病毒
	②鼻病毒	②鼻病毒
	③普通冠状病毒	③呼吸道合胞病毒
60 岁及以上	①流感病毒	①呼吸道合胞病毒
	②鼻病毒	②流感病毒
	③普通冠状病毒	③鼻病毒

图 1 哨点医院门急诊流感样病例数占门急诊就诊总数比例每周变化趋势

图 2 哨点医院门急诊流感样病例呼吸道样本病原体核酸检测阳性率每周变化趋势

图 3 哨点医院住院严重急性呼吸道感染病例呼吸道样本病原体核酸检测阳性率每周变化趋势